

REMARKS ON A PAPER BY JUHÁSZ AND KUNEN

SAKAÉ FUCHINO AND STEFAN GESCHKE

ABSTRACT. We give an equivalent, but simpler formulation of the axiom SEP introduced by Juhász and Kunen in [8]. Our formulation shows that many of the consequences of the weak Freese-Nation Property of $\mathcal{P}(\omega)$ studied in [4] already follow from SEP. We show that it is consistent that SEP holds while $\mathcal{P}(\omega)$ fails to have the (\aleph_1, \aleph_0) -ideal property introduced in [2]. This answers a question addressed independently by Fuchino and by Kunen. We also consider some natural variants of SEP and show that certain changes in the definition of SEP do not lead to a different principle. This answers a question addressed by Blass.

1. INTRODUCTION

In [8], Juhász and Kunen introduced a new principle, SEP, which is designed to capture some consequences of CH and holds in Cohen models, i.e., models of set theory obtained by adding Cohen reals to models of CH. The principle says that for sufficiently large χ there are many good elementary submodels M of the structure (\mathcal{H}_χ, \in) such that $\mathcal{P}(\omega) \cap M$ is nicely embedded in $\mathcal{P}(\omega)$.

Different notions of *good* and *many* elementary submodels and *nice* embeddings lead to different principles. In this paper we study the relation between SEP and the (\aleph_1, \aleph_0) -ideal property of Dow and Hart [2], which is defined by the same pattern. The corresponding embeddings are *sep*-embeddings and σ -embeddings.

Definition 1.1. Let A and B be Boolean algebras such that $A \leq B$. Then $A \leq_{sep} B$ if and only if for every $b \in B$ and every uncountable set $T \subseteq A \restriction b$ there is $a \in A \restriction b$ such that $\{c \in T : c \leq a\}$ is uncountable. Let $A \leq_\sigma B$ if and only if for every $b \in B$, $A \restriction b$ has a countable cofinal subset.

Observe that $A \leq_\sigma B$ implies $A \leq_{sep} B$.

The good elementary submodels will be the same for both SEP and the (\aleph_1, \aleph_0) -ideal property: later we show that \leq_{sep} can be replaced by \leq_σ in the following definition of SEP. The real difference between SEP and the (\aleph_1, \aleph_0) -ideal property lies in the interpretation of *many*.

Extending the notions defined by Juhász and Kunen, we regard SEP and the (\aleph_1, \aleph_0) -ideal property as properties of general Boolean algebras, not only of $\mathcal{P}(\omega)$.

Date: September 12, 2003.

2000 Mathematics Subject Classification. Primary: 03E35; Secondary: 03E05, 03E17, 03E65.

Key words and phrases. weak Freese-Nation property, SEP, Cohen model, almost disjoint number. The first author thanks Hiroshi Sakai for pointing out a gap in the proof of an earlier version of Theorem 8.14.

Definition 1.2. For a cardinal χ let \mathcal{M}_χ be the set of elementary submodels M of \mathcal{H}_χ such that $|M| = \aleph_1$ and $[M]^{\aleph_0} \cap M$ is cofinal in $[M]^{\aleph_0}$. For a Boolean algebra A , $\text{SEP}(A)$ is the statement “for all sufficiently large regular χ there are cofinally many $M \in \mathcal{M}_\chi$ such that $A \cap M \leq_{\text{sep}} A$ ”. The axiom SEP introduced in [8] is $\text{SEP}(\mathcal{P}(\omega))$.

A has the (\aleph_1, \aleph_0) -ideal property (IDP) if and only if for all sufficiently large regular χ and all $M \in \mathcal{M}_\chi$ with $A \in M$, $A \cap M \leq_\sigma A$ holds. We write $\text{IDP}(A)$ if A has the IDP.

As usual, we identify an algebraic structure $(A, f_1, \dots, f_n, R_1, \dots, R_m)$ with its underlying set A . It should be made clear that by “ $A \in M$ ” we really mean $(A, f_1, \dots, f_n, R_1, \dots, R_m) \in M$. The role of the cardinal χ is analyzed in Section 8, where we calculate precisely when a regular cardinal is “sufficiently large” in the definition of SEP .

Since \leq_σ is stronger than \leq_{sep} , for a Boolean algebra A , $\text{IDP}(A)$ implies $\text{SEP}(A)$. In Section 2 we show that the relation \leq_{sep} in the definition of SEP can be replaced by \leq_σ , i.e., SEP and IDP are very similar. In Section 8 we observe that it does not make a difference in the definition of SEP if we replace “there are cofinally many $M \in \mathcal{M}_\chi$ ” by “there is $M \in \mathcal{M}_\chi$ ” or by “there are stationarily many (in $[\mathcal{H}_\chi]^{\aleph_1}$) $M \in \mathcal{M}_\chi$ ”. This shows the importance of the results of Section 6 and Section 7, namely that SEP is really weaker than IDP .

If the universe is not very complex, that is, if some very weak version of the \square -principle together with $\text{cf}([\mu]^{\aleph_0}) = \mu^+$ holds for all singular cardinals μ of countable cofinality, or if the Boolean algebras under consideration are small, then the (\aleph_1, \aleph_0) -ideal property is equivalent to the weak Freese-Nation property studied in [6]:

Definition 1.3. A Boolean algebra A has the weak Freese-Nation property (WFN) if and only if there is a function $f : A \rightarrow [A]^{\leq \aleph_0}$ such that for all $a, b \in A$ with $a \leq b$ there is $c \in f(a) \cap f(b)$ with $a \leq c \leq b$. f is called a WFN-function for A . We write $\text{WFN}(A)$ for “ A has the WFN”.

It is easy to check that if f is a WFN-function for A and $B \leq A$ is closed under f , then $B \leq_\sigma A$. In [6] Fuchino, Koppelberg, and Shelah characterized the WFN using elementary submodels and σ -embeddings. They showed

Theorem 1.4. *A Boolean algebra A has the WFN if and only if for all sufficiently large χ and all $M \preccurlyeq \mathcal{H}_\chi$ with $\aleph_1 \subseteq M$ and $A \in M$, $A \cap M \leq_\sigma A$.*

Since $\aleph_1 \subseteq M$ for all $M \in \mathcal{M}_\chi$, it is clear that for a Boolean algebra A , $\text{IDP}(A)$ follows from $\text{WFN}(A)$. As mentioned above, IDP and WFN are equivalent in many cases. Some of these cases are captured by the following lemma, which follows from the results in [7].

Lemma 1.5. *If A is a Boolean algebra of size $< \aleph_\omega$ or if 0^\sharp does not exist, then $\text{IDP}(A)$ holds if and only if $\text{WFN}(A)$ does.*

The formulation using 0^\sharp is chosen here just for simplicity. As mentioned above, what is really needed is only a certain very weak assumption at singular cardinals

of countable cofinality. On the other hand, it is known that the lemma does not hold without any such additional assumption (see [7] or [5]).

In [4] many interesting consequences of $\text{WFN}(\mathcal{P}(\omega))$ have been found. Concerning the combinatorics of the reals, a universe satisfying $\text{WFN}(\mathcal{P}(\omega))$ behaves very similar to a Cohen model. In particular, the values of the popular cardinal invariants of the continuum, that is, those studied in [1], have the same values in a model with $\text{WFN}(\mathcal{P}(\omega))$ as in a Cohen model with the same size of the continuum.

Our characterization of SEP in terms of \leq_σ rather than \leq_{sep} shows that the axiom $\text{SEP}(\mathcal{P}(\omega))$ is sufficient to determine at least some of the smaller cardinal invariants of the continuum.

Juhász and Kunen already proved that another consequence of $\text{WFN}(\mathcal{P}(\omega))$, the principle $C_2^s(\omega_2)$ introduced in [9], follows from the weaker assumption $\text{SEP}(\mathcal{P}(\omega))$.

2. SEP IS SIMILAR TO IDP

As mentioned above, \leq_σ implies \leq_{sep} . If the subalgebras under consideration are of size $\leq \aleph_1$, then the two relations are in fact the same.

Lemma 2.1. *Let A and B be Boolean algebras with $A \leq_{sep} B$ and $|A| = \aleph_1$. Then $A \leq_\sigma B$.*

Proof. Let $b \in B$ and assume for a contradiction that $A \restriction b$ is not countably generated. Let $(a_\alpha)_{\alpha < \omega_1}$ enumerate $A \restriction b$. By recursion on $\alpha < \omega_1$ define a sequence $(c_\alpha)_{\alpha < \omega_1}$ in $A \restriction b$ such that for all $\alpha < \omega_1$, $a_\alpha \leq c_\alpha$ and c_α is not in the ideal of A generated by $\{c_\beta : \beta < \alpha\}$.

Now let $T = \{c_\alpha : \alpha < \omega_1\}$. We claim that T is a counterexample to $A \leq_{sep} B$. For let $a \in T$. Then there is $\beta < \omega_1$ with $a = a_\beta$. Thus $a_\beta \leq c_\beta$. By the construction of the sequence $(c_\alpha)_{\alpha < \omega_1}$, there are only countably many elements of T below c_β . Hence, there are only countably many elements of T below a , contradicting $A \leq_{sep} B$. \square

Thus, we have

Corollary 2.2. *For every Boolean algebra A , $\text{SEP}(A)$ holds if and only if for all sufficiently large regular χ there are cofinally many $M \in \mathcal{M}_\chi$ with $A \cap M \leq_\sigma A$.*

In this characterization, it is easily seen that many interesting consequences of $\text{WFN}(\mathcal{P}(\omega))$ already follow from $\text{SEP}(\mathcal{P}(\omega))$. In the proofs of most of the results in [4] it is only used that under $\text{WFN}(\mathcal{P}(\omega))$, for some sufficiently large χ there are cofinally many $M \in \mathcal{M}_\chi$ with $\mathcal{P}(\omega) \cap M \leq_\sigma \mathcal{P}(\omega)$. The following theorem collects some of the consequences of $\text{SEP}(\mathcal{P}(\omega))$ that follow from the arguments given in [4].

A subset of $[\omega]^{\aleph_0}$ is called groupwise dense if it is closed under taking almost subsets and non-meager with respect to the topology on $[\omega]^{\aleph_0}$ inherited from 2^ω when identifying $[\omega]^{\aleph_0}$ with a subset of 2^ω .

Theorem 2.3. *Assume $\text{SEP}(\mathcal{P}(\omega))$. Then the following cardinal invariants of the continuum are \aleph_1 .*

1. $\text{non}(\mathcal{M})$, the smallest size of a non-meager subset of \mathbb{R} and
2. \mathfrak{g} , the smallest size of a family of groupwise dense subsets of $[\omega]^{\aleph_0}$ with empty intersection.

Moreover, if CH fails, then $\text{cov}(\mathcal{M})$, the minimal size of a family of non-meager subsets of \mathbb{R} covering \mathbb{R} , is at least \aleph_2 .

It was also proved in [4] that $\text{WFN}(\mathcal{P}(\omega))$ implies that \mathfrak{a} , the smallest size of a maximal almost disjoint family in $\mathcal{P}(\omega)$, is \aleph_1 . In the proof it is sufficient to assume $\text{IDP}(\mathcal{P}(\omega))$. The situation with \mathfrak{a} under $\text{SEP}(\mathcal{P}(\omega))$ is more subtle and will be discussed in section 8.

For more information about these cardinal invariants see e.g. [1].

3. $\text{SEP}(\mathcal{P}(\omega))$ HOLDS IN COHEN MODELS

In [5] it was shown that $\text{WFN}(\mathcal{P}(\omega))$ can fail in a Cohen model, assuming the consistency of some very large cardinal. In [7] it was shown that large cardinal assumptions are necessary for this.

$\text{SEP}(\mathcal{P}(\omega))$ however, is always true in a Cohen model. We include a proof of this fact (Theorem 3.1). It follows that $\text{WFN}(\mathcal{P}(\omega))$ does not follow from $\text{SEP}(\mathcal{P}(\omega))$ in ZFC (assuming the consistency of certain large cardinals). We do not know whether $\text{IDP}(\mathcal{P}(\omega))$ is always true in a Cohen model.

In section 7 we shall show, without large cardinal assumptions, that $\text{SEP}(\mathcal{P}(\omega))$ does not imply $\text{IDP}(\mathcal{P}(\omega))$.

Theorem 3.1. *Let V be a model of CH and suppose that G is $\text{Fn}(\kappa, 2)$ -generic over V . Then $V[G] \models \text{SEP}(\mathcal{P}(\omega))$.*

The proof of this theorem relies on the following series of lemmas. The first lemma was proved in [10].

Lemma 3.2. *Let M be a transitive model of set theory and let $x \subseteq \omega$ be a Cohen real over M . Then $\mathcal{P}(\omega) \cap M \leq_\sigma \mathcal{P}(\omega) \cap M[x]$.*

Lemma 3.3. *Let P be a c.c.c. partial order and let $M \in \mathcal{M}_\chi$ be such that $P \in M$. Let G be P -generic over the ground model V . Then $M[G] \in \mathcal{M}_\chi^{V[G]}$.*

Proof. Let M and G be as above. Then $M[G] \preceq \mathcal{H}_\chi[G]$ by c.c.c. of P . Let $X \subseteq M[G]$ be countable. In $V[G]$, there is a countable set $C \subseteq M$ of P -names such that $X \subseteq \{\dot{x}_G : \dot{x} \in C\}$. Again by the c.c.c. of P , we may assume that $C \in V$ and C is countable in V . By $M \in \mathcal{M}_\chi^V$, we may assume $C \in M$. Since M contains a name for G , $G \in M[G]$ and thus $X \subseteq \{\dot{x}_G : \dot{x} \in C\} \in M[G]$. \square

Lemma 3.4. *Assume CH. Then $M \in \mathcal{M}_\chi$ implies that $[M]^{\leq \aleph_0} \subseteq M$.*

In particular, for all $M \in \mathcal{M}_\chi$, if $P \subseteq M$ is a c.c.c. partial order, then every nice P -name for a subset of ω is contained in M .

Proof. Suppose that $x \in [M]^{\aleph_0}$. By $M \in \mathcal{M}_\chi$ there is $y \in [M]^{\aleph_0} \cap M$ such that $x \subseteq y$. By CH there is a surjection $f : \omega_1 \rightarrow \mathcal{P}(y)$. By elementarity, there is such f

in M . Let $\alpha < \omega_1$ be such that $f(\alpha) = x$. Since $\omega_1 \subseteq M$, we have $\alpha \in M$ and thus $x = f(\alpha) \in M$.

Now let $P \subseteq M$ be a c.c.c. partial order. Since P is c.c.c., every nice P -name for a subset of ω is a countable subset of $\{\check{n} : n \in \omega\} \times P \subseteq M$. By the first part of the lemma, every such set is an element of M . \square

Lemma 3.5. *Let V and G be as in Theorem 3.1. Let $\chi \in V$ be large enough and let $M \in (\mathcal{M}_\chi)^V$ be such that $\kappa \in M$. Then $\mathcal{P}(\omega) \cap M[G] = \mathcal{P}(\omega) \cap V[G \cap \text{Fn}(\kappa \cap M, 2)]$.*

Proof. First let $x \in \mathcal{P}(\omega) \cap M[G]$. In M there is a name \dot{x} for x which is a nice name for a subset of ω . Since $\text{Fn}(\kappa, 2)$ satisfies the c.c.c., in V there is a countable set $X \subseteq \kappa$ such that \dot{x} is an $\text{Fn}(X, 2)$ -name. We can find such an X in M . Since $\aleph_0 \subseteq M$, $X \subseteq M$. Therefore \dot{x} is an $\text{Fn}(\kappa \cap M, 2)$ -name and thus $x \in V[G \cap \text{Fn}(\kappa \cap M, 2)]$.

For the converse let $x \in V[G \cap \text{Fn}(\kappa \cap M, 2)]$. Pick a nice $\text{Fn}(\kappa \cap M, 2)$ -name \dot{x} for x . Clearly, $\text{Fn}(\kappa \cap M, 2) \subseteq M$. Therefore Lemma 3.4 applies, and we get $\dot{x} \in M$. This shows $x \in M[G]$. \square

Proof of Theorem 3.1. We argue in $V[G]$. Let χ be sufficiently large and let $M \in \mathcal{M}_\chi$. It is sufficient to show that there is $M' \in \mathcal{M}_\chi$ with $M \subseteq M'$ and $\mathcal{P}(\omega) \cap M' \leq_\sigma \mathcal{P}(\omega)$.

Since $\text{Fn}(\kappa, 2)$ has the c.c.c., in V there is a set X of size \aleph_1 of $\text{Fn}(\kappa, 2)$ -names such that every element of M has a name in X . Let $N \in \mathcal{M}_\chi^V$ be such that $X \subseteq N$. Clearly, $M \subseteq N[G]$. By Lemma 3.3, $N[G] \in \mathcal{M}_\chi$. By Lemma 3.5, $\mathcal{P}(\omega) \cap N[G] = \mathcal{P}(\omega) \cap V[G \cap \text{Fn}(\kappa \cap N, 2)]$. Since $V[G]$ is an $\text{Fn}(\kappa \setminus N, 2)$ -generic extension over $V[G \restriction \text{Fn}(\kappa \cap N, 2)]$, it follows from Lemma 3.2 that $\mathcal{P}(\omega) \cap N[G] \leq_\sigma \mathcal{P}(\omega)$. This shows that, in $V[G]$, the set of $M' \in \mathcal{M}_\chi$ with $\mathcal{P}(\omega) \cap M' \leq_\sigma \mathcal{P}(\omega)$ is cofinal in $[\mathcal{H}_\chi]^{\aleph_1}$. \square

4. IDP FOR PARTIAL ORDERS

In [6] the WFN has been defined for partial orders, not only for Boolean algebras. In this section and the next, we do the same for SEP and IDP. We have to liberalize our definition of \leq_σ .

Definition 4.1. Let P and Q be partial orders with $Q \leq P$, i.e., $Q \subseteq P$ and the orders on Q and P agree on Q . For $p \in P$ let $Q \restriction p = \{q \in Q : q \leq p\}$ and $Q \upharpoonright p = \{q \in Q : q \geq p\}$. Now $Q \leq_\sigma P$ if and only if for all $p \in P$, $Q \restriction p$ has a countable cofinal subset and $Q \upharpoonright p$ has a countable coinital subset.

It is clear that for Boolean algebras A and B , if A is a subalgebra of B , then $A \leq_\sigma B$ holds in the Boolean algebraic sense if and only if $A \leq_\sigma B$ holds for the partial orders. Now we can extend the notions IDP, SEP, and WFN to partial orders.

Definition 4.2. For a partial order P , $\text{SEP}(P)$ holds if and only if for all sufficiently large regular χ there are cofinally many $M \in \mathcal{M}_\chi$ with $P \cap M \leq_\sigma P$.

IDP(P) holds if and only if for all sufficiently large regular χ and for all $M \in \mathcal{M}_\chi$ with $P \in M$, $P \cap M \leq_\sigma P$.

WFN(P) holds if and only if for all sufficiently large regular χ and all $M \preceq \mathcal{H}_\chi$ with $\aleph_1 \subseteq M$ and $P \in M$, $P \cap M \leq_\sigma P$.

It is easy to check that the extended versions of SEP, IDP, and WFN agree with the old ones on Boolean algebras. For SEP this relies on Corollary 2.2, for WFN on Theorem 1.4.

Let us look at the partial orders $([\kappa]^{\leq \aleph_0}, \subseteq)$. Assuming the consistency of some large cardinal, it was shown in [7] that $\text{WFN}([\aleph_\omega]^{\leq \aleph_0})$ does not follow from GCH. However, we have

Theorem 4.3. *CH implies IDP($[\kappa]^{\leq \aleph_0}$) for every cardinal κ .*

Proof. Assume CH and let χ be sufficiently large and regular. Let $M \in \mathcal{M}_\chi$ be such that $[\kappa]^{\leq \aleph_0} \in M$. Then we have $\kappa \in M$. By Lemma 3.4, $M \cap [\kappa]^{\leq \aleph_0} = [\kappa \cap M]^{\leq \aleph_0}$.

Thus it remains to show that $[\kappa \cap M]^{\leq \aleph_0} \leq_\sigma [\kappa]^{\leq \aleph_0}$. Suppose $x \in [\kappa]^{\leq \aleph_0}$. If $x \not\subseteq \kappa \cap M$, then $[\kappa \cap M]^{\leq \aleph_0} \upharpoonright x = \emptyset$. Thus, $[\kappa \cap M]^{\leq \aleph_0} \upharpoonright x$ is either empty or has a minimal element, namely x .

$[\kappa \cap M]^{\leq \aleph_0} \upharpoonright x$ always has a maximal element, namely $x \cap M$. This finishes the proof of the theorem. \square

This theorem can be regarded as a parallel of Theorem 3.1. It follows that, assuming the consistency of some large cardinal, it is consistent that there is a partial order that has the IDP, but not the WFN.

5. COMPLETE BOOLEAN ALGEBRAS SATISFYING SEP

Just as the WFN, SEP is hereditary with respect to order retracts. A partial order P is an order retract of a partial order Q if there are order preserving maps $e : P \rightarrow Q$ and $f : Q \rightarrow P$ such that $f \circ e = \text{id}_P$. If P and Q are Boolean algebras and e and f are (Boolean) homomorphisms, then we call P a retract of Q .

Lemma 5.1. *Let P and Q be partial orders such that P is an order retract of Q . Then $\text{SEP}(Q)$ implies $\text{SEP}(P)$.*

Proof. Let χ be large enough and $M \preceq \mathcal{H}_\chi$. Suppose $M \cap Q \leq_\sigma Q$ and $P, Q \in M$. We show $P \cap M \leq_\sigma P$.

Since M knows that P is a retract of Q , M contains order preserving maps $e : P \rightarrow Q$ and $f : Q \rightarrow P$ such that $f \circ e = \text{id}_P$. Let $p \in P$. Since $Q \cap M \leq_\sigma Q$, there is a countable set $C \subseteq Q \cap M$ such that C is cofinal in $Q \cap M \upharpoonright e(p)$.

Claim. $f[C]$ is cofinal in $P \cap M \upharpoonright p$.

Let $q \in P \cap M \upharpoonright p$. Since $e(q) \in Q \cap M \upharpoonright e(p)$, there is $c \in C$ such that $e(q) \leq c \leq e(p)$. Now $q \leq f(c) \leq p$, which proves the claim.

By the same argument, $P \cap M \upharpoonright p$ has a countable coinitial subset. This implies $\text{SEP}(P)$. \square

If P is a complete lattice and P embeds into Q via e , then there is a map $f : Q \rightarrow P$ with $f \circ e = \text{id}_P$, namely the map defined by $f(q) = \sup\{p \in P : e(p) \leq q\}$ for all $q \in Q$. Thus, a complete lattice which embeds into a partial order Q with $\text{SEP}(Q)$ also has the property SEP .

Note that if A is a complete Boolean algebra and A embeds into B in the Boolean algebraic sense, then A is a retract of B by Sikorski's extension criterion. If A is an infinite complete Boolean algebra, then A has a maximal antichain of size \aleph_0 and $\mathcal{P}(\omega)$ embeds into A , again by Sikorski's extension criterion. Since $\mathcal{P}(\omega)$ is complete, it is a retract of A .

It follows that $\text{SEP}(\mathcal{P}(\omega))$ holds if there is any infinite complete Boolean algebra A with $\text{SEP}(A)$. Note that for this it is not necessary to use Sikorski's criterion; our statement about complete lattices suffices.

While $\text{SEP}(\mathcal{P}(\omega))$ is consistent with (but not a theorem of) ZFC, $\text{SEP}(\mathcal{P}(\omega_1))$ fails. We prove this in a series of lemmas. Together with Lemma 5.1 this will imply that all complete Boolean algebras A with $\text{SEP}(A)$ satisfy the c.c.c. It should be pointed out that the proof of $\neg \text{WFN}(\mathcal{P}(\omega_1))$ given in [6] also works for SEP . However, we believe that our argument is simpler.

In the following χ always denotes a sufficiently large regular cardinal.

Lemma 5.2. a) $\neg \text{SEP}(\omega_2 + 1)$

b) If $\text{SEP}(A)$ holds, then A does not have a chain of order type ω_2 .

Proof. For a) note that for each $M \in \mathcal{M}_\chi$, $\omega_2 \cap M$ is an ordinal of cofinality \aleph_1 . It follows that for each $\alpha \in \omega_2 \setminus M$, $(\omega_2 + 1) \cap M \upharpoonright \alpha$ has uncountable cofinality. In particular, $(\omega_2 + 1) \cap M \not\leq_\sigma \omega_2 + 1$.

For b) let A be a Boolean algebra such that ω_2 embeds into A . Clearly, $\omega_2 + 1$ also embeds into A . But $\omega_2 + 1$ is a complete lattice. Thus $\omega_2 + 1$ is an order retract of A . Now $\neg \text{SEP}(A)$ follows from Lemma 5.1 together with part a). \square

Lemma 5.3. $\neg \text{SEP}(\mathcal{P}(\omega_1)/[\omega_1]^{\leq \aleph_0})$

Proof. By Lemma 5.2, it suffices to show that $\mathcal{P}(\omega_1)/[\omega_1]^{\leq \aleph_0}$ has a chain of order-type ω_2 .

For $f, g \in \omega_1^{\omega_1}$ let $f <^* g$ if and only if $\{\alpha < \omega_1 : f(\alpha) \geq g(\alpha)\}$ is countable. If $(f_\gamma)_{\gamma < \omega_2}$ is a $<^*$ -increasing sequence in $\omega_1^{\omega_1}$, then $(\{(\alpha, \beta) \in \omega_1 \times \omega_1 : \beta \leq f_\gamma(\alpha)\})_{\gamma < \omega_2}$ gives rise to a strictly increasing sequence in $\mathcal{P}(\omega_1 \times \omega_1)/[\omega_1 \times \omega_1]^{\leq \aleph_0}$ of order type ω_2 . It is thus sufficient to construct a $<^*$ -increasing sequence of order type ω_2 in $\omega_1^{\omega_1}$. But this is easy using the natural diagonalization argument to get an $<^*$ -upper bound for any set $F \subseteq \omega_1^{\omega_1}$ of size $\leq \aleph_1$. \square

Lemma 5.4. $\neg \text{SEP}(\mathcal{P}(\omega_1))$

Proof. Let $M \in \mathcal{M}_\chi$. Suppose $\mathcal{P}(\omega_1) \cap M \leq_\sigma \mathcal{P}(\omega_1)$. We show that

$$(\mathcal{P}(\omega_1)/[\omega_1]^{\leq \aleph_0}) \cap M \leq_\sigma \mathcal{P}(\omega_1)/[\omega_1]^{\leq \aleph_0}.$$

This suffices for the lemma since together with Lemma 5.3 it implies that there are not cofinally many $N \in \mathcal{M}_\chi$ with $\mathcal{P}(\omega_1) \cap N \leq_\sigma \mathcal{P}(\omega_1)$. Note that $(\mathcal{P}(\omega_1) \cap$

$M)/([\omega_1]^{\leq \aleph_0} \cap M)$ is essentially the same as $(\mathcal{P}(\omega_1)/[\omega_1]^{\leq \aleph_0}) \cap M$ since M includes a cofinal subset of $[\omega_1]^{\leq \aleph_0}$, namely the countable ordinals.

Let $a \in \mathcal{P}(\omega_1)$. Let C be a cofinal subset of $\mathcal{P}(\omega_1) \cap M \restriction a$. Let D be the set of all classes modulo $[\omega_1]^{\leq \aleph_0}$ of elements of C and let \bar{a} be the class of a modulo $[\omega_1]^{\leq \aleph_0}$.

Claim. D is cofinal in $((\mathcal{P}(\omega_1)/[\omega_1]^{\leq \aleph_0}) \cap M) \restriction \bar{a}$.

Let $b \in \mathcal{P}(\omega_1) \cap M$ be such that $b \setminus a$ is countable. Let $\alpha < \omega_1$ be such that $b \setminus \alpha \subseteq a$. Since $\alpha \in M$, $b \setminus \alpha \in M$. Therefore, there is $c \in C$ such that $b \setminus \alpha \subseteq c \subseteq a$. The claim clearly follows from this. This finishes the proof of the lemma. \square

Let A be a complete Boolean algebra not satisfying the c.c.c. Since A is complete, A has an maximal antichain of size \aleph_1 . This antichain gives rise to an embedding of the algebra of finite-cofinite subsets of ω_1 into A . Since A is complete, this embedding extends to all of $\mathcal{P}(\omega_1)$ by Sikorski's extension criterion. Since $\mathcal{P}(\omega_1)$ is complete, it follows that $\mathcal{P}(\omega_1)$ is a retract of A . Using Lemma 5.1 this gives

Corollary 5.5. *Let A be a complete Boolean algebra. If $\text{SEP}(A)$ holds, then A satisfies the c.c.c.*

6. AN EXAMPLE IN ZFC

In this section we show in ZFC that there is a Boolean algebra A which satisfies $\text{SEP}(A)$ but not $\text{IDP}(A)$.

Definition 6.1. Let $E_{\aleph_1}^{\aleph_2} = \{\alpha < \omega_2 : \text{cf}(\alpha) = \aleph_1\}$. For all $\alpha \in E_{\aleph_1}^{\aleph_2}$ fix an increasing sequence $(\delta_\beta^\alpha)_{\beta < \omega_1}$ which is cofinal in α and consists of successor ordinals. For $S \subseteq E_{\aleph_1}^{\aleph_2}$ let A^S be the Boolean algebra defined as follows: Let $A_0^S = \{0, 1\}$. Suppose $\alpha < \omega_2$ is a limit ordinal and A_β^S has already been defined for all $\beta < \alpha$. Let $A_\alpha^S = \bigcup_{\beta < \alpha} A_\beta^S$. Now suppose A_α^S has been defined and $\alpha \in \omega_2 \setminus S$. Let $A_{\alpha+1}^S = A_\alpha^S(x_\alpha)$ where x_α is independent over A_α^S . Suppose that A_α^S has already been defined and $\alpha \in S$. Let $A_{\alpha+1}^S = A_\alpha^S(x_\alpha)$ where $x_\alpha \notin A_\alpha^S$, $A_\alpha^S \restriction x_\alpha$ is generated by $\{x_{\delta_\beta^\alpha} : \beta < \omega_1\}$ and $A_\alpha^S \restriction -x_\alpha$ is $\{0\}$. Finally let $A^S = \bigcup_{\alpha < \omega_2} A_\alpha^S$.

Fix a sufficiently large regular cardinal χ . Recall that for every $M \in \mathcal{M}_\chi$, $\omega_1 \subseteq M$ and thus, $M \cap \omega_2 \in \omega_2$.

Lemma 6.2. *Let $X \subseteq \mathcal{H}_\chi$ be of size \aleph_1 . Then $C_X = \{M \cap \omega_2 : M \in \mathcal{M}_\chi \wedge X \subseteq M\}$ includes an \aleph_1 -club of ω_2 , that is, an unbounded set which is closed under limits of subsets of cofinality \aleph_1 .*

Proof. By recursion, we define an increasing sequence $(M_\alpha)_{\alpha < \omega_2}$ in \mathcal{M}_χ such that

- (i) $X \subseteq M_0$,
- (ii) for all $\alpha < \omega_2$, $\alpha \in M_\alpha$, and
- (iii) if $\alpha < \omega_2$ is a limit ordinal of cofinality \aleph_1 , then $M_\alpha = \bigcup_{\beta < \alpha} M_\beta$.

This construction can be carried out since \mathcal{M}_χ is cofinal in $[\mathcal{H}_\chi]^{\aleph_1}$ and closed under unions of chains of length ω_1 . Let $C = \{\omega_2 \cap M_\alpha : \alpha < \omega_2\}$. Then C is unbounded in ω_2 by (ii) and closed under limits of subsets of cofinality \aleph_1 by (iii). By (i), $C \subseteq C_X$. \square

From Lemma 6.2 we get

Lemma 6.3. *Let S be a stationary subset of ω_2 such that $S \subseteq E_{\aleph_1}^{\aleph_2}$. Then for cofinally many $M \in \mathcal{M}_\chi$ we have $\omega_2 \cap M \in S$.*

Proof. Let $X \in [\mathcal{H}_\chi]^{\aleph_1}$. By Lemma 6.2, the set C_X includes an \aleph_1 -club C of ω_2 . Let \overline{C} be the closure of C , i.e., C together with all limit points of C . Then \overline{C} is club in ω_2 and $C = \overline{C} \cap E_{\aleph_1}^{\aleph_2}$. Since S is stationary and a subset of $E_{\aleph_1}^{\aleph_2}$, $\overline{C} \cap S = C \cap S$ is non-empty and thus, there is $M \in \mathcal{M}_\chi$ such that $X \subseteq M$ and $M \cap \omega_2 \in S$. \square

Using Lemma 6.3, we can show that for a suitably chosen set S , the Boolean algebra A^S constructed above satisfies SEP but not IDP.

Theorem 6.4. *There is a Boolean algebra A with $\text{SEP}(A)$ but not $\text{IDP}(A)$.*

Proof. Fix two disjoint stationary subsets S_0 and S_1 of ω_2 with $S_0 \cup S_1 = E_{\aleph_1}^{\aleph_2}$. Let $A = A^{S_1}$.

Claim 1: $\text{SEP}(A)$

Let $M \in \mathcal{M}_\chi$ be such that $\alpha = M \cap \omega_2 \in S_0$ and $A \in M$. Then $A \cap M = A_\alpha^{S_1}$. Since $\alpha \notin S_1$ and by the construction of A^{S_1} , $A_\alpha^{S_1} \leq_\sigma A$. This proves the claim since there are cofinally many $M \in \mathcal{M}_\chi$ with $M \cap \omega_2 \in S_0$ and $A \in M$ by Lemma 6.3.

Claim 2: $\neg \text{IDP}(A)$

By Lemma 6.3, there is $M \in \mathcal{M}_\chi$ such that $\alpha = M \cap \omega_2 \in S_1$ and $A \in M$. As above, $A \cap M = A_\alpha^{S_1}$. By the construction of A^{S_1} , $A_\alpha^{S_1} \not\leq_\sigma A$. In other words, M witnesses the failure of $\text{IDP}(A)$. \square

7. $\text{SEP}(\mathcal{P}(\omega))$ DOES NOT IMPLY $\text{IDP}(\mathcal{P}(\omega))$

In this section we use the idea of the proof of Theorem 6.4 to construct a model of set theory where $\text{SEP}(\mathcal{P}(\omega))$ holds while $\text{IDP}(\mathcal{P}(\omega))$ fails.

Theorem 7.1. *It is consistent that $\text{SEP}(\mathcal{P}(\omega))$ holds but $\text{IDP}(\mathcal{P}(\omega))$ fails.*

Proof. Suppose the ground model V satisfies CH and let S_0 and S_1 be as in the proof of Theorem 6.4. For each $\alpha \in E_{\aleph_1}^{\aleph_2}$ let $(\delta_\beta^\alpha)_{\beta < \omega_1}$ be as in Definition 6.1.

Our strategy is to perform a finite support iteration of c.c.c. forcings over V of length ω_2 where we add only Cohen reals most of the time. However, at stage $\alpha \in S_1$ we add a new subset x_α of ω such that the Cohen reals added at the stages δ_β^α , $\beta < \omega_1$, are almost contained in x_α . Note that we consider the Cohen reals to be subsets of ω . This construction should be viewed as the forcing version of the construction in the proof of Theorem 6.4.

We now define the iteration $(P_\alpha, Q_\alpha)_{\alpha < \omega_2}$. The underlying sets of the Q_α 's will be absolute, but not the orders. Thus we will not define each Q_α as a P_α -name but as the underlying set of Q_α in V , also named Q_α , together with a P_α -name $\dot{\leq}_\alpha$ for the order on Q_α .

For all $\alpha \notin S_1$ let Q_α be Cohen forcing, i.e., $\text{Fn}(\omega, 2)$. Let $\dot{\leq}_\alpha$ be the canonical name for the usual order on $\text{Fn}(\omega, 2)$, i.e., reverse inclusion. For $\alpha \in S_1$ and $\beta < \omega_1$

let $\dot{x}_{\delta_\beta^\alpha}$ be a $P_{\delta_\beta^\alpha+1}$ -name for the Cohen real added by $Q_{\delta_\beta^\alpha}$. Set

$$Q_\alpha = \{(f, F) : f \in {}^{<\omega}2, F \in [\{\delta_\beta^\alpha : \beta \in \omega_1\}]^{<\aleph_0}\}$$

and let $\dot{\leq}_\alpha$ be a name for a relation \leq on Q_α such that $(f, F) \leq (f', F')$ if and only if $f' \subseteq f$, $F' \subseteq F$, and for all $\delta \in F'$, if $n \in \dot{x}_\delta \cap \text{dom}(f \setminus f')$, then $f(n) = 1$.

As usual, for each $\alpha < \omega_2$ let P_α be the finite support iteration of the Q_β , $\beta < \alpha$, where each $Q_\alpha = (Q_\alpha, \dot{\leq}_\alpha)$ is considered as a P_α -name for the appropriate partial order. Let P be the direct limit of the P_α , $\alpha < \omega_2$. For convenience, by the absoluteness of the elements of the Q_α 's, we may assume that the elements of each P_α and of P are elements of $\prod_{\beta < \omega_2} Q_\beta$ with finite support. For each condition $p \in P$ let $\text{supt}(p)$ be its support.

Note that P is c.c.c. since the Q_α 's are σ -centered. Let G be P -generic over V . For $\alpha < \omega_2$ let $G_\alpha = G \cap P_\alpha$.

The easier part of the proof of the theorem is to show that in $V[G]$, $\text{IDP}(\mathcal{P}(\omega))$ fails. To see this, we need

Lemma 7.2. *For $\alpha \in S_1$, $V[G] \models \mathcal{P}(\omega) \cap V[G_\alpha] \not\leq_\sigma \mathcal{P}(\omega)$.*

Proof. We argue in $V[G]$. Let $\alpha \in S_1$. Note that $\mathcal{P}(\omega) \cap V[G_\alpha] \leq_\sigma \mathcal{P}(\omega)$ if and only if $(\mathcal{P}(\omega) \cap V[G_\alpha]) / \text{fin} \leq_\sigma \mathcal{P}(\omega) / \text{fin}$.

For $y \in \mathcal{P}(\omega)$ let \bar{y} be the equivalence class of y modulo fin . Let x be the subset of ω generically added by Q_α .

Claim. $(\mathcal{P}(\omega) \cap V[G_\alpha]) / \text{fin} \restriction \bar{x}$ is generated by the classes modulo fin of the Cohen reals $\{x_{\delta_\beta^\alpha} : \beta < \omega_1\}$ added by the $Q_{\delta_\beta^\alpha}$'s. In particular, $(\mathcal{P}(\omega) \cap V[G_\alpha]) / \text{fin} \not\leq_\sigma \mathcal{P}(\omega) / \text{fin}$.

It follows from the construction of Q_α that for all $\beta < \omega_1$, $\bar{x}_{\delta_\beta^\alpha} \leq \bar{x}$. Let $a \in \mathcal{P}(\omega) \cap V[G_\alpha]$. Suppose that a is not almost included in the union of a finite subset of $\{x_{\delta_\beta^\alpha} : \beta < \omega_1\}$. Then for every $n \in \omega$ the set of conditions in Q_α which force that there is $m \geq n$ such that $m \in a$ but $m \notin x$ is easily seen to be dense in Q_α . It follows that a is not almost included in x . This shows the claim and finishes the proof of Lemma 7.2. \square

By Lemma 6.3, in V there is $M \in \mathcal{M}_\chi$ containing P such that $\alpha = M \cap \omega_2 \in S_1$. Now $M[G] \cap \omega_2 = \alpha$ since P is c.c.c. If $\dot{x} \in M$ is a P -name for a subset of ω , then M also contains a nice P -name \dot{y} for the same subset of ω . By c.c.c., \dot{y} only uses countably many conditions from P . These conditions are already contained in P_β for some $\beta < \alpha$. Since V satisfies CH and since $P_\beta \in M$ (and thus $P_\beta \subseteq M$), $\dot{y} \in M$ by Lemma 3.4. It follows that $\mathcal{P}(\omega) \cap M[G] = \mathcal{P}(\omega) \cap V[G_\alpha]$. Therefore, $M[G]$ shows that $\mathcal{P}(\omega)$ does not satisfy IDP in $V[G]$.

To see that $\text{SEP}(\mathcal{P}(\omega))$ holds in $V[G]$ we need

Lemma 7.3. *If $\alpha < \omega_2$ and $\alpha \notin S_1$, then $V[G] \models \mathcal{P}(\omega) \cap V[G_\alpha] \leq_\sigma \mathcal{P}(\omega)$.*

Proof. By Lemma 3.2, it is sufficient to show that for $\alpha \in \omega_2 \setminus S_1$ every real in $V[G] \setminus V[G_\alpha]$ is contained in a Cohen extension of $V[G_\alpha]$.

Let $\alpha \in \omega_2 \setminus S_1$. Let $x \in \mathcal{P}(\omega)$, but $x \notin V[G_\alpha]$. Then there is an P -name $\dot{x} \in V$ for x . By c.c.c., we may assume that \dot{x} uses only countably many conditions from P . Our plan is to find $P_{\dot{x}} \subseteq P$ such that

- (1) $P_\alpha \subseteq P_{\dot{x}}$ and P_α is completely embedded in $P_{\dot{x}}$,
- (2) \dot{x} is an $P_{\dot{x}}$ -name,
- (3) $P_{\dot{x}}$ is completely embedded in P , and
- (4) the quotient $P_{\dot{x}} : G_\alpha$ is equivalent to $\text{Fn}(\omega, 2)$.

This suffices for the lemma. For suppose $P_{\dot{x}}$ is as above. It is not hard to see that (3) implies that $P_{\dot{x}} : G_\alpha$ is completely embedded in $P : G_\alpha$. Note that by (1), it is reasonable to consider $P_{\dot{x}} : G_\alpha$. By (2), \dot{x} can be regarded as a $P_{\dot{x}} : G_\alpha$ -name. Thus by (4), x is contained in a Cohen extension of $V[G_\alpha]$ and the lemma follows.

It remains to construct $P_{\dot{x}}$ and to show that it has the required properties. For a condition $p \in P$ let

$$\overline{\text{supt}}(p) = \text{supt}(p) \cup \bigcup \{F : \exists \beta \in \text{supt}(p) \cap S_1 \exists f(p(\beta) = (f, F))\}.$$

Let

$$X = \alpha \cup \bigcup \{\text{supt}(p) : \dot{x} \text{ uses the condition } p\}$$

and $P_{\dot{x}} = \{p \in P : \overline{\text{supt}}(p) \subseteq X\}$.

Claim. $P_{\dot{x}}$ has the properties (1)–(4).

(1) $P_\alpha \subseteq P_{\dot{x}}$ follows from the definitions. $P_{\dot{x}}$ can be viewed as finite support iteration which has P_α as an initial segment. Thus P_α is completely embedded in $P_{\dot{x}}$.

(2) It follows from the definitions that \dot{x} is an $P_{\dot{x}}$ -name.

(3) We have to show the following:

- (i) $\forall p, q \in P_{\dot{x}} (p \perp_{P_{\dot{x}}} q \Rightarrow p \perp_P q)$,
- (ii) $\forall p \in P \exists q \in P_{\dot{x}} \forall r \in P_{\dot{x}} (r \leq q \Rightarrow r \not\perp_P p)$.

For (i) observe that for all $p, q \in P$ with $p \not\perp_P q$ there is $r \in P$ such that $r \leq p, q$ and $\overline{\text{supt}}(r) \subseteq \overline{\text{supt}}(p) \cup \overline{\text{supt}}(q)$. Therefore, if $p, q \in P_{\dot{x}}$ are compatible in P , then they are in $P_{\dot{x}}$.

(ii) is what really requires work. Let $p \in P$. Let $q' \in P_{\dot{x}}$ be the condition with support $\text{supt}(p) \cap X$ such that for all $\beta \in (\text{supt}(p) \cap X) \setminus S_1$, $q'(\beta) = p(\beta)$ and for all $\beta \in \text{supt}(p) \cap X \cap S_1$, $q'(\beta) = (f, F \cap X)$ where f and F are such that $p(\beta) = (f, F)$. q' does not yet work for q in (ii). We have to extend it a little.

Let q be the condition with the same support as q' such that $q(\beta) = q'(\beta)$ for all $\beta \in \text{supt}(q') \setminus S_1$. Now fix $\beta \in \text{supt}(q') \cap S_1$. Let f and F be such that $q'(\beta) = (f, F)$. Let $m \in \omega$ be such that for all $\gamma \in \text{supt}(p) \setminus S_1$, $\text{dom}(p(\gamma)) \subseteq m$. Let $g \in 2^{<\omega}$ be such that $m \subseteq \text{dom}(g)$, $f \subseteq g$, and $g(n) = 1$ for all $n \in \text{dom}(g) \setminus \text{dom}(f)$. Now set $q(\beta) = (g, F)$.

Subclaim. q works for (ii).

Let $r \in P_{\dot{x}}$ be such that $r \leq q$. We have to construct a common extension $s \in P$ of p and r . As above, we build an approximation s' of s first. For $\beta \in S_1$ with $p(\beta) = (f, F)$ and $r(\beta) = (f', F')$ let $s'(\beta) = (f \cup f', F \cup F')$. Note that $f \cup f'$ is a

function since by the definition of q and by $r \leq q$, we even have $f \subseteq f'$ whenever $\beta \in \text{supt}(r)$. Note that this definition makes sense if $\beta \notin \text{supt}(r) \cap \text{supt}(p)$ since the largest element of Q_β is simply (\emptyset, \emptyset) (for $\beta \in S_1$).

For $\beta \in \omega_2 \setminus S_1$ let $s'(\beta) = p(\beta) \cup r(\beta)$. Again, $p(\beta) \cup r(\beta)$ is a function since for $\beta \in \text{supt}(r)$, $p(\beta) \subseteq r(\beta)$ by $r \leq q$ and the definition of q . It is easy to see that s' extends r .

It may happen that $s' \not\leq p$. However, we can extend s' to a condition $s \leq p$ by adding some Cohen conditions (deciding more of the Cohen reals involved). Let $s(\beta) = s'(\beta)$ for all $\beta \in S_1$. For $\beta \in S_1$ we have to make sure that $s \restriction \beta$ forces $s(\beta)$ to be below $p(\beta)$.

For all $\beta \in S_1$ let f_β , F_β , f'_β , and F'_β be such that $s(\beta) = (f_\beta, F_\beta)$ and $p(\beta) = (f'_\beta, F'_\beta)$. Then for all $\beta \in S_1$ we have $f'_\beta \subseteq f_\beta$ and $F'_\beta \subseteq F_\beta$. For all $\delta \in F'_\beta$ we want to have $\delta \in \text{supt}(s)$ and $s(\delta) \Vdash \forall n \in \text{dom}(f_\beta) \setminus \text{dom}(f'_\beta) (f_\beta(n) \geq \dot{x}_\delta(n))$. This can be accomplished. Just let z be a sufficiently long finite sequence of zeros and put $s(\delta) = s'(\delta) \frown z$ for every $\delta \in \bigcup \{F_\beta : \beta \in \text{supt}(s') \cap S_1\}$. Note that there are only finitely many δ 's to be considered. For every $\beta \in \omega_2$ for which $s(\beta)$ has not yet been defined let $s(\beta) = s'(\beta)$.

It is straight forward to check that s is a common extension of r and p . This completes the proof of the subclaim and thus shows that $P_{\dot{x}}$ is completely embedded in P .

(4) Note that any two elements of $P_{\dot{x}}$ that agree on $[\alpha, \omega_2)$ are equivalent in $P_{\dot{x}} : G_\alpha$, i.e., they will be identified in the completion of $P_{\dot{x}} : G_\alpha$. But since $\{\delta_\gamma^\beta : \gamma \in \omega_1\} \cap \alpha$ is countable for all $\beta \in [\alpha, \omega_2) \cap S_1$ and $X \setminus \alpha$ is countable, there are only countably many possibilities for $p \restriction [\alpha, \omega_2)$ for $p \in P_{\dot{x}}$. Therefore, the completion of $P_{\dot{x}} : G_\alpha$ has a countable dense subset. Since below each element of $P_{\dot{x}} : G_\alpha$ there are two incompatible elements (in $P_{\dot{x}} : G_\alpha$), $P_{\dot{x}} : G_\alpha$ is equivalent to $\text{Fn}(\omega, 2)$. This finishes the proof of the claim and of the lemma. \square

Since there are cofinally many $M \in (\mathcal{M}_\chi)^V$ with $M \cap \omega_2 \in S_0$ by Lemma 6.3, the set $\{M[G] : M \in \mathcal{M}_\chi \wedge M \cap \omega_2 \in S_0\}$ is cofinal in $(\mathcal{M}_\chi)^{V[G]}$. As above, for all $M \in (\mathcal{M}_\chi)^V$, $\alpha = M[G] \cap \omega_2 = M \cap \omega_2$ and $\mathcal{P}(\omega) \cap M[G] = \mathcal{P}(\omega) \cap V[G_\alpha]$. Now it follows from Lemma 7.3 that $\text{SEP}(\mathcal{P}(\omega))$ holds in $V[G]$. This finishes the proof of the theorem. \square

8. VARIANTS OF SEP

It is tempting to define a new class of partial orders by replacing “cofinally many $M \in \mathcal{H}_\chi$ ” in the definition of SEP by “stationarily many $M \in \mathcal{H}_\chi$ ”. However, the class of partial orders with this modified notion of SEP coincides with the class of partial orders with the original SEP. Also, one arrives at the same notion if “there are cofinally $M \in \mathcal{H}_\chi$ ” is weakened to “there is $M \in \mathcal{H}_\chi$ ”.

For a partial order P and a regular cardinal χ such that $P \in \mathcal{H}_\chi$ let

$$\mathcal{M}(P, \chi) = \{M \in \mathcal{M}_\chi : P \in M \wedge P \cap M \leq_\sigma P\}.$$

Theorem 8.1. *Let P be any partial order. Then for $\kappa = \max(|\text{trcl}(P)|, \aleph_1)$ the following are equivalent:*

- (1) $\text{SEP}(P)$
- (2) *There is a regular cardinal $\chi > \kappa$ such that $\mathcal{M}(P, \chi)$ is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$.*
- (3) *For every regular cardinal $\chi > \kappa$, $\mathcal{M}(P, \chi)$ is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$.*
- (4) *There is a regular cardinal $\chi > 2^\kappa$ such that $\mathcal{M}(P, \chi)$ is non-empty.*
- (5) *For every regular cardinal $\chi > 2^\kappa$, $\mathcal{M}(P, \chi)$ is non-empty.*

Clearly, (1) follows from (3) and implies (4). The remaining part of Theorem 8.1 is a special case of the following lemma, which does not have to do anything with σ -embeddings. For a set A , a family $\mathcal{F} \subseteq \mathcal{P}(A)$, and a regular cardinal χ with $A, \mathcal{F} \in \mathcal{H}_\chi$ let

$$\mathcal{M}(A, \mathcal{F}, \chi) = \{M \in \mathcal{M}_\chi : A, \mathcal{F} \in M \wedge A \cap M \in \mathcal{F}\}.$$

For a partial order P , $\mathcal{M}(P, \chi)$ is simply $\mathcal{M}(P, \{Q \subseteq P : Q \leq_\sigma P\}, \chi)$.

Lemma 8.2. *Let A be a set and $\mathcal{F} \subseteq \mathcal{P}(A)$. Then for $\kappa = \max(|\text{trcl}(A)|, \aleph_1)$ the following are equivalent:*

- (1) *There is a regular cardinal $\chi \geq \kappa^+$ such that $\mathcal{M}(A, \mathcal{F}, \chi)$ is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$.*
- (2) *For every regular cardinal $\chi \geq \kappa^+$, $\mathcal{M}(A, \mathcal{F}, \chi)$ is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$.*
- (3) *There is a regular cardinal $\chi > 2^\kappa$ such that $\mathcal{M}(A, \mathcal{F}, \chi)$ is non-empty.*
- (4) *For every regular cardinal $\chi > 2^\kappa$, $\mathcal{M}(A, \mathcal{F}, \chi)$ is non-empty.*

The reason for considering κ^+ and 2^κ in the formulation of this lemma is that κ^+ is the least cardinal $\chi > \aleph_1$ with $A \in \mathcal{H}_\chi$ and the size of \mathcal{H}_{κ^+} is 2^κ . The proof of Lemma 8.2 uses two arguments: one for stepping up in cardinality and one for stepping down. We start with decreasing cardinals. Fix A, \mathcal{F} , and κ as in Lemma 8.2.

Lemma 8.3. *Let $\chi, \mu > \kappa$ be regular cardinals with $2^{<\chi} < \mu$. If $\mathcal{M}(A, \mathcal{F}, \mu)$ is non-empty, then $\mathcal{M}(A, \mathcal{F}, \chi)$ is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$.*

Proof. Suppose that $\mathcal{M}(A, \mathcal{F}, \chi)$ is not stationary in $[\mathcal{H}_\chi]^{\aleph_1}$. We may assume that $\chi > \kappa$ is minimal with this property. Let $M \in \mathcal{M}(A, \mathcal{F}, \mu)$.

Since $|\mathcal{H}_\chi|^{\aleph_1} = 2^{<\chi} < \mu$, we have $[\mathcal{H}_\chi]^{\aleph_1}, \mathcal{M}(A, \mathcal{F}, \chi) \in \mathcal{H}_\mu$ and χ is definable in \mathcal{H}_μ with the parameters A and \mathcal{F} . Therefore, $\chi \in M$ and M knows that $\mathcal{M}(A, \mathcal{F}, \chi)$ is not stationary in $[\mathcal{H}_\chi]^{\aleph_1}$.

It follows that M contains a club \mathcal{C} of $[\mathcal{H}_\chi]^{\aleph_1}$ which is disjoint from $\mathcal{M}(A, \mathcal{F}, \chi)$. By elementarity, $M \cap \mathcal{H}_\chi \subseteq \bigcup(\mathcal{C} \cap M)$. Since $\aleph_1 \subseteq M$, $\bigcup(\mathcal{C} \cap M) \subseteq M \cap \mathcal{H}_\chi$. Since $M \cap [M]^{\aleph_0}$ is cofinal in $[M]^{\aleph_0}$, $\mathcal{C} \cap M$ is countably directed. It follows that $\bigcup(M \cap \mathcal{C})$ is the union of an increasing chain of length ω_1 of elements of \mathcal{C} . Therefore, $M \cap \mathcal{H}_\chi = \bigcup(\mathcal{C} \cap M) \in \mathcal{C}$. It is easily checked that $M \cap \mathcal{H}_\chi \in \mathcal{M}_\chi$.

Since $A \subseteq \mathcal{H}_\chi$, $A \cap M \cap \mathcal{H}_\chi = A \cap M \in \mathcal{F}$. Thus, $M \cap \mathcal{H}_\chi \in \mathcal{C} \cap \mathcal{M}(A, \mathcal{F}, \chi)$, contradicting the choice of \mathcal{C} . \square

Lemma 8.4. *For all regular cardinals $\chi, \mu > \kappa$ with $\chi < \mu$, if $\mathcal{M}(A, \mathcal{F}, \chi)$ is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$, then $\mathcal{M}(A, \mathcal{F}, \mu)$ is non-empty.*

Proof. We use a refined Skolem hull operator to find $\tilde{M} \in \mathcal{M}_\mu$ with $A, \mathcal{F} \in \tilde{M}$ and $\tilde{M} \cap A \in \mathcal{F}$. Fix a well-ordering \sqsubset on \mathcal{H}_μ . For $\alpha < \omega_1$ let sk_α denote the Skolem hull operator on \mathcal{H}_μ with respect to the built-in Skolem functions of the structure $(\mathcal{H}_\mu, \in, \sqsubset, A, \mathcal{F}, \text{sk}_\beta)_{\beta < \alpha}$ where A and \mathcal{F} are considered as constants. For $X \subseteq \mathcal{H}_\chi$ let

$$\text{sk}^*(X) = \bigcup \{ \text{sk}_\alpha(Y) : Y \in X \wedge |Y| \leq \aleph_0 \wedge \alpha < \omega_1 \}.$$

Claim. Let $X \in [\mathcal{H}_\mu]^{\aleph_1}$ be such that $X \cap [X]^{\leq \aleph_0}$ is cofinal in $[X]^{\leq \aleph_0}$. Then

- (i) $X \subseteq \text{sk}^*(X)$ and $A, \mathcal{F} \in \text{sk}^*(X)$,
- (ii) $|\text{sk}^*(X)| = \aleph_1$,
- (iii) $[\text{sk}^*(X)]^{\leq \aleph_0} \cap \text{sk}^*(X)$ is cofinal in $[\text{sk}^*(X)]^{\leq \aleph_0}$, and
- (iv) $\text{sk}^*(X) \prec \mathcal{H}_\mu$.

Moreover,

- (v) for all $X, Y \subseteq \mathcal{H}_\mu$ with $X \subseteq Y$, $\text{sk}^*(X) \subseteq \text{sk}^*(Y)$ and
- (vi) if $(X_\alpha)_{\alpha < \delta}$ is an increasing sequence of subsets of \mathcal{H}_μ and $X = \bigcup_{\alpha < \delta} X_\alpha$, then $\text{sk}^*(X) = \bigcup_{\alpha < \delta} \text{sk}^*(X_\alpha)$.

For (i) let $x \in X$. By our assumptions on X , there is $Y \in X \cap [X]^{\leq \aleph_0}$ such that $\{x\} \subseteq Y$. Now $x \in Y \subseteq \text{sk}_0(Y)$ and thus $x \in \text{sk}^*(X)$. This shows $X \subseteq \text{sk}^*(X)$. $A, \mathcal{F} \in \text{sk}^*(X)$ since $A, \mathcal{F} \in \text{sk}_0(Y)$ for every countable element Y of X .

Statement (ii) follows from the fact that for every countable set $Y \subseteq \mathcal{H}_\mu$ and every $\alpha < \omega_1$, $\text{sk}_\alpha(Y)$ is again countable.

For (iii) let Y be a countable subset of $\text{sk}^*(X)$. For every $n \in \omega$ fix $\alpha_n < \omega_1$ and a countable set $Y_n \in X$ such that $Y \subseteq \bigcup_{n \in \omega} \text{sk}_{\alpha_n}(Y_n)$. By our assumptions on X , there is $Z \in X \cap [X]^{\leq \aleph_0}$ such that $\{Y_n : n \in \omega\} \subseteq Z$. Let $\beta = \sup_{n \in \omega} (\alpha_n + 1)$. Now for every $n \in \omega$, $\text{sk}_{\alpha_n}(Y_n) \in \text{sk}_\beta(Z)$ by the choice of sk_β . Since $\text{sk}_\beta(Z)$ is an elementary submodel of \mathcal{H}_μ and since the $\text{sk}_{\alpha_n}(Y_n)$ are countable, for every $n \in \omega$ we also have $\text{sk}_{\alpha_n}(Y_n) \subseteq \text{sk}_\beta(Z)$. It follows that $Y \subseteq \text{sk}_\beta(Z)$. Clearly, $\text{sk}_\beta(Z)$ is a countable subset of $\text{sk}^*(X)$. We are done with the proof of (iii) if we can show $\text{sk}_\beta(Z) \in \text{sk}^*(X)$. But this is easy. Just let $Z' \in X \cap [X]^{\leq \aleph_0}$ be such that $Z \in Z'$. Now $\text{sk}_\beta(Z) \in \text{sk}_{\beta+1}(Z') \subseteq \text{sk}^*(X)$.

For (iv) it suffices to show that for all finite subsets F of $\text{sk}^*(X)$ there is an elementary submodel M of \mathcal{H}_μ such that $F \subseteq M \subseteq \text{sk}^*(X)$. Let F be a finite subset of $\text{sk}^*(X)$. As before, there are $\beta < \omega_1$ and $Z \in X \cap [X]^{\leq \aleph_0}$ such that $F \subseteq \text{sk}_\beta(Z)$. By the definition of sk^* , $\text{sk}_\beta(Z) \subseteq \text{sk}^*(X)$. And $\text{sk}_\beta(Z)$ is an elementary submodel of \mathcal{H}_μ .

Statements (v) and (vi) follow immediately from the definition of sk^* . This finishes the proof of the claim.

Now consider the set

$$\mathcal{C} = \{M \in [\mathcal{H}_\chi]^{\aleph_1} : \text{sk}^*(M) \cap \mathcal{H}_\chi = M\}.$$

From the properties of sk^* it follows that \mathcal{C} is club in $[\mathcal{H}_\chi]^{\aleph_1}$. Since $\mathcal{M}(X, \mathcal{F}, \chi)$ is stationary, there is $M \in \mathcal{C} \cap \mathcal{M}(X, \mathcal{F}, \chi)$. Let $\tilde{M} = \text{sk}^*(M)$. By the properties of sk^* , $\tilde{M} \in \mathcal{M}_\chi$ and $A, \mathcal{F} \in \tilde{M}$. Moreover, $\tilde{M} \cap X = M \cap X \in \mathcal{F}$. In other words, $\tilde{M} \in \mathcal{M}(X, \mathcal{F}, \mu)$. \square

Proof of Lemma 8.2. We start from (3). Suppose there is a regular cardinal $\chi > 2^\kappa$ such that $\mathcal{M}(X, \mathcal{F}, \chi)$ is non-empty. Then, by Lemma 8.3, $\mathcal{M}(X, \mathcal{F}, \kappa^+)$ is stationary in $[\mathcal{H}_{\kappa^+}]^{\aleph_1}$. This implies (1).

Now suppose that (1) holds. Then there is a regular cardinal $\chi > \kappa$ such that $\mathcal{M}(X, \mathcal{F}, \chi)$ is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$. By Lemma 8.4, there are arbitrarily large regular cardinals μ such that $\mathcal{M}(X, \mathcal{F}, \mu)$ is non-empty. By Lemma 8.3, this implies that $\mathcal{M}(X, \mathcal{F}, \mu)$ is stationary for every regular $\mu > \kappa$, i.e., (2) holds. The implications (2) \Rightarrow (4) and (4) \Rightarrow (3) are trivial. \square

At the moment, we do not know whether $\mathfrak{a} = \aleph_1$ follows from $\text{SEP}(\mathcal{P}(\omega))$. However, we can show that a variant of $\text{SEP}(\mathcal{P}(\omega))$ which is called $\text{SEP}^{+-}(\mathcal{P}(\omega))$ here (see below) implies $\mathfrak{a} = \aleph_1$.

In the following let χ always denote a regular cardinal.

Definition 8.5.

$$\mathcal{M}_\chi^\sqsubset = \{M : M \prec \mathcal{H}_\chi, |M| = \aleph_1, \text{ and there is a well-ordering } \sqsubset \text{ on } M \text{ of order type } \omega_1 \text{ such that for every } a \in M, \sqsubset \cap (M_{\sqsubset a})^2 \in M\}$$

where $M_{\sqsubset a} = \{x \in M : x \sqsubset a\}$.

Definition 8.6. Let P be a partial order.

(1) $\text{SEP}^+(P)$ if $\{M \in \mathcal{M}_\chi^\sqsubset : P \cap M \leq_\sigma P\}$ is cofinal in $[\mathcal{H}_\chi]^{\aleph_1}$ for every sufficiently large χ .

(2) $\text{SEP}^{+-}(P)$ if $\{M \in \mathcal{M}_\chi^\sqsubset : P \cap M \leq_\sigma P\}$ is non-empty for a sufficiently large χ .

For A, \mathcal{F}, χ as in the definition of $\mathcal{M}(A, \mathcal{F}, \chi)$, let

$$\mathcal{M}^\sqsubset(A, \mathcal{F}, \chi) = \{M \in \mathcal{M}_\chi^\sqsubset : A, \mathcal{F} \in M \wedge A \cap M \in \mathcal{F}\}.$$

Then it is easy to see that Lemma 8.3 with $\mathcal{M}^\sqsubset(A, \mathcal{F}, \chi)$ in place of $\mathcal{M}(A, \mathcal{F}, \chi)$ also holds. As in the proof of Lemma 8.2, we obtain the following equivalence:

$$\text{SEP}^+(P)$$

$$\Leftrightarrow \{M \in \mathcal{M}_\chi^\sqsubset : P \cap M \leq_\sigma P\} \text{ is stationary for every sufficiently large } \chi$$

$$\Leftrightarrow \{M \in \mathcal{M}_\chi^\sqsubset : P \cap M \leq_\sigma P\} \text{ is non-empty for every sufficiently large } \chi$$

Lemma 8.7. $\mathcal{M}_\chi^\sqsubset \subseteq \mathcal{M}_\chi$.

Proof. Suppose that $M \in \mathcal{M}_\chi^\sqsubset$ and \sqsubset is a well-ordering of M as in the definition of $\mathcal{M}_\chi^\sqsubset$. For $X \in [M]^{\aleph_0}$ let $x \in M$ be such that $X \subseteq M_{\sqsubset x}$. Then $|M_{\sqsubset x}| \leq \aleph_0$ and $M_{\sqsubset x} \in M$. This shows that $[M]^{\aleph_0} \cap M$ is cofinal in $[M]^{\aleph_0}$. Hence $M \in \mathcal{M}_\chi$. \square

Lemma 8.8. Suppose that $M \in \mathcal{M}_\chi^\sqsubset$. Then M is internally approachable in the sense of [3], i.e.,

- (*) M is the union of a continuously increasing chain of countable elementary submodels $(M_\alpha)_{\alpha < \omega_1}$ of M such that $(M_\beta)_{\beta \leq \alpha} \in M_{\alpha+1}$ for all $\alpha < \omega_1$.

Proof. Suppose that $M \in \mathcal{M}_\chi^\square$ and \sqsubset is a well-ordering of M as in the definition of \mathcal{M}_χ^\square .

Let $x_\alpha \in M$, $\alpha < \omega_1$ be defined inductively such that

- (0) $M_{\sqsubset x_\alpha}$ is an elementary submodel of (M, \in, \sqsubset) ;
- (1) If α is a limit ordinal then x_α is the limit of x_β , $\beta < \alpha$;
- (2) If α is a successor, say $\alpha = \beta + 1$, then x_α is minimal with respect to \sqsubset such that $x_\beta \sqsubset x_\alpha$, $\sqsubset \cap (M_{\sqsubset x_\beta})^2 \in M_{\sqsubset x_\alpha}$ and (0).

Note that, in (2), the construction is possible since $\sqsubset \cap (M_{\sqsubset x_\beta})^2 \in M$ by the definition of $M \in \mathcal{M}_\chi^\square$. By (0) and since $(x_\beta)_{\beta \leq \alpha}$ is definable in M with the parameter $\sqsubset \cap (M_{\sqsubset x_\alpha})^2$ we have $(x_\beta)_{\beta \leq \alpha} \in M_{\sqsubset x_{\alpha+1}}$. It follows that $(M_{\sqsubset x_\beta})_{\beta \leq \alpha} \in M_{\sqsubset x_{\alpha+1}}$. Also $M = \bigcup_{\alpha < \omega_1} M_{\sqsubset x_\alpha}$ since \sqsubset has order type ω_1 . Thus $(M_{\sqsubset x_\alpha})_{\alpha < \omega_1}$ is a sequence as required in (*). \square

The property (*) in Lemma 8.8 almost characterizes elements of \mathcal{M}_χ^\square :

Lemma 8.9. *Let $<^*$ be a well-ordering of \mathcal{H}_χ of order type $|\mathcal{H}_\chi|$. If $(M_\alpha)_{\alpha < \omega_1}$ is a continuously increasing sequence of countable elementary submodels of $(\mathcal{H}_\chi, \in, <^*)$ such that $(M_\beta)_{\beta \leq \alpha} \in M_{\alpha+1}$ for all $\alpha < \omega_1$, then $M = \bigcup_{\alpha < \omega_1} M_\alpha$ is an element of \mathcal{M}_χ^\square .*

Proof. For each $x \in M$, let $\alpha_x = \min\{\alpha < \omega_1 : x \in M_{\alpha+1}\}$. Let \sqsubset be the linear ordering on M defined by

$$x \sqsubset y \Leftrightarrow \alpha_x < \alpha_y \vee (\alpha_x = \alpha_y \wedge x <^* y).$$

Clearly \sqsubset is a well-ordering on M . \sqsubset has order type ω_1 since every initial segment of M with respect to \sqsubset is countable and M itself is uncountable.

Let $x \in M$. We show that $\sqsubset \cap (M_{\sqsubset x})^2 \in M$. Let $\alpha^* = \alpha_x + 2$. In M_{α^*} , $\sqsubset \cap (M_{\sqsubset x})^2$ is definable from $<^* \cap (M_{\alpha_x+1})^2$ and $(M_\beta)_{\beta \leq \alpha_x+1}$. By elementarity and since $<^* \cap (M_{\alpha_x+1})^2$ and $(M_\beta)_{\beta \leq \alpha_x+1}$ are elements of M_{α^*} , it follows that $\sqsubset \cap (M_{\sqsubset x})^2 \in M_{\alpha^*} \subseteq M$. \square

By Lemma 8.8 and Lemma 8.9, there are club many $M \prec \mathcal{H}_\chi$ of size \aleph_1 such that $M \in \mathcal{M}_\chi^\square$ if and only if M is internally approachable — namely those M with $M \prec (\mathcal{H}_\chi, \in, <^*)$ for some fixed $<^*$ as above.

Lemma 8.10. \mathcal{M}_χ^\square is stationary in $[\mathcal{H}_\chi]^{\aleph_1}$.

Proof. Suppose that $\mathcal{C} \subseteq [\mathcal{H}_\chi]^{\aleph_1}$ is closed unbounded. We show that $\mathcal{M}_\chi^\square \cap \mathcal{C} \neq \emptyset$.

Let $<^*$ be a well-ordering of \mathcal{H}_χ of order type $|\mathcal{H}_\chi|$. Let $(M_\alpha)_{\alpha < \omega_1}$ be a continuously increasing chain of countable elementary submodels of $(\mathcal{H}_\chi, \mathcal{C}, \in, <^*)$ such that $(M_\beta)_{\beta \leq \alpha} \in M_{\alpha+1}$ for all $\alpha < \omega_1$. Let $M = \bigcup_{\alpha < \omega_1} M_\alpha$. Then $M \in \mathcal{M}_\chi^\square$ by Lemma 8.9. Since $\omega_1 \subseteq M$, we have $N \subseteq M$ for all $N \in \mathcal{C} \cap M$. By elementarity $\mathcal{C} \cap M$ is a directed system and $M = \bigcup(\mathcal{C} \cap M)$. Since $|M| = \aleph_1$, it follows that $M \in \mathcal{C}$. \square

Lemma 8.11. *For a partial order P*

- (1) $\text{IDP}(P)$ *implies* $\text{SEP}^+(P)$;
- (2) $\text{SEP}^+(P)$ *implies* $\text{SEP}^{+-}(P)$;
- (3) $\text{SEP}^{+-}(P)$ *implies* $\text{SEP}(P)$;

$$\text{WFN}(P) \Rightarrow \text{IDP}(P) \Rightarrow \text{SEP}^+(P) \Rightarrow \text{SEP}^{+-}(P) \Rightarrow \text{SEP}(P)$$

Proof. (1) follows from Lemma 8.7. (2) is clear from definitions. (3) follows from Theorem 8.1. \square

Theorem 8.12. *Assume $\text{SEP}^{+-}(\mathcal{P}(\omega))$. Then $\mathfrak{a} = \aleph_1$.*

Proof. Let χ be sufficiently large and $M^* \in \mathcal{M}_\chi^\square$ be such that $\mathcal{P}(\omega) \cap M^* \leq_\sigma \mathcal{P}(\omega)$. Since $|M^*| = \aleph_1$, it is enough to show that there is a MAD-family $\subseteq M^*$.

Let \sqsubset be a well-ordering of M^* as in the definition of \mathcal{M}_χ^\square . By Lemma 8.8, there is an increasing sequence $(M_\alpha)_{\alpha < \omega_1}$ of countable elementary submodels of M^* such that $\bigcup_{\alpha < \omega_1} M_\alpha = M$ and $(M_\beta)_{\beta \leq \alpha} \in M_{\alpha+1}$ for all $\alpha < \omega_1$.

Let $(a_\alpha)_{\alpha < \omega_1}$ be such that

- (1) $\{a_n : n \in \omega\}$ is a partition of ω with $(a_n)_{n \in \omega} \in M_0$;
- (2) For $\alpha \geq \omega$, $a_\alpha \in [\omega]^{\aleph_0} \cap M_{\alpha+1}$ is minimal (with respect to \sqsubset) with the following property:
 - (α) a_α is almost disjoint to each a_β , $\beta < \alpha$.
 - (β) $\forall x \in [\omega]^{\aleph_0} \cap M_\alpha \left(\forall u \in [\alpha]^{<\aleph_0} (|x \setminus \bigcup_{\beta \in u} a_\beta| = \aleph_0) \rightarrow |a_\alpha \cap x| = \aleph_0 \right)$.

By (1) and (2), $(a_\beta)_{\beta < \alpha}$ is definable in $M_{\alpha+1}$ using the parameters $\sqsubset \cap (M_\alpha)^2$, $(M_\beta)_{\beta \leq \alpha} \in M_{\alpha+1}$. Hence $(a_\beta)_{\beta < \alpha} \in M_{\alpha+1}$ and so the inductive construction of a_α satisfying (α) and (β) can be accomplished in $M_{\alpha+1}$.

By (1) and (2)(α), $\{a_\beta : \beta < \omega_1\}$ is pairwise almost disjoint. To show that it is maximal, suppose that it were not. Then there is some $b \in [\omega]^{\aleph_0}$ such that b is almost disjoint to all of a_α 's. Let $\{b_n : n \in \omega\} \subseteq \mathcal{P}(\omega) \cap M^*$ be a countable generator of $(\mathcal{P}(\omega) \cap M^*) \upharpoonright b$. Let $\alpha^* < \omega_1$ be such that $\{b_n : n \in \omega\} \subseteq M_{\alpha^*}$. Since a_{α^*} and b are almost disjoint there is $n^* \in \omega$ such that $|b_{n^*} \cap a_{\alpha^*}| < \aleph_0$. By (2)(β), there is a $u \in [\alpha]^<\aleph_0$ such that $|b_{n^*} \setminus \bigcup_{\beta \in u} a_\beta| < \aleph_0$. As $b \subseteq b_{n^*}$, it follows that $|b \setminus \bigcup_{\beta \in u} a_\beta| < \aleph_0$. But this is a contradiction to the choice of b . \square

Irreversibility of the implications in Lemma 8.11 cannot be proved in ZFC:

Lemma 8.13. *Assume CH. Then $\mathcal{M}_\chi = \mathcal{M}_\chi^\square$. In particular, for every partial order P , $\text{SEP}(P)$ if and only if $\text{SEP}^+(P)$.*

Proof. $\mathcal{M}_\chi^\square \subseteq \mathcal{M}_\chi$ by Lemma 8.7. To show $\mathcal{M}_\chi \subseteq \mathcal{M}_\chi^\square$, suppose $M \in \mathcal{M}_\chi$. Then by Lemma 3.4 $[M]^{\aleph_0} \subseteq M$. Let \sqsubset be an arbitrary well-ordering of M of order type ω_1 . Then, for every $x \in M$, $M_{\sqsubset x}$ and $\sqsubset \cap (M_{\sqsubset x})^2$ are countable subsets of M and hence, by CH, elements of M . This shows that $M \in \mathcal{M}_\chi^\square$. \square

Even under $\neg\text{CH}$, SEP and SEP^+ can be equivalent for partial orders with an “ \aleph_2 -version” of IDP.

Let us say that a partial order P has the \aleph_2 -IDP if for any sufficiently large χ and $M \prec \mathcal{H}_\chi$, if $|M| = \aleph_2$, $P \in M$ and $[M]^{\aleph_1} \cap M$ is cofinal in $[M]^{\aleph_1}$ then $P \cap M \leq_{\aleph_2} P$ where $P \leq_{\aleph_2} Q$ is defined just as in Definition 4.1 with “countable” there replaced by “of cardinality $< \aleph_2$ ”.

Note that every partial order of cardinality $\leq \aleph_2$ has the \aleph_2 -IDP.

Theorem 8.14. *Assume \square_{ω_1} . For any partial order P with the \aleph_2 -IDP, $\text{SEP}(P)$ if and only if $\text{SEP}^+(P)$.*

For the proof of Theorem 8.14 we use the fact that the class of partial orders with SEP is closed under \leq_σ -suborders.

Lemma 8.15. *For partial orders P and Q , if $\text{SEP}(P)$ and $Q \leq_\sigma P$ then $\text{SEP}(Q)$.*

Proof. Fix a sufficiently large regular χ . It is enough to show that, for every $M \prec \mathcal{H}_\chi$ with $P, Q \in M$, if $P \cap M \leq_\sigma P$ then $Q \cap M \leq_\sigma Q$. To see this, let $x_0 \in Q$ and we show that $Q \cap M \restriction x_0$ has a countable cofinal subset. (That $Q \cap M \restriction x_0$ has a countable cointial subset can be proved similarly.) By $P \cap M \leq_\sigma P$ there is a countable set $X \subseteq (P \cap M) \restriction x_0$ such that X is cofinal in $(P \cap M) \restriction x_0$. By $M \models Q \leq_\sigma P$ and elementarity, for every $x \in X$ we can find $X_x \in M$ such that $M \models “X_x \text{ is cofinal in } Q \restriction x”$. Then $X_x \subseteq M$ and X_x is a countable cofinal subset of $(Q \cap M) \restriction x$. Letting $Y = \bigcup_{x \in X} X_x$, we have $Y \subseteq (Q \cap M) \restriction x_0$ and Y is countable. Y is a cofinal subset of $(Q \cap M) \restriction x_0$: If $y \in (Q \cap M) \restriction x_0$, then in particular $y \in (P \cap M) \restriction x_0$. Hence there is $x \in X$ such that $y \leq x$. Since $M \models y \in Q \restriction x$, there is $x' \in X_x \subseteq Y$ such that $y \leq x'$. \square

Note that the proof above actually shows that each of the variants of SEP considered above (and also IDP and WFN) is closed under \leq_σ -suborders.

Proof of Theorem 8.14. If $|P| < \aleph_2$, then the assertion of the theorem is trivial. Hence we may assume $|P| \geq \aleph_2$. If $\text{SEP}^+(P)$ then $\text{SEP}(P)$ by Lemma 8.11 (2). So we assume $\text{SEP}(P)$ and prove $\text{SEP}^+(P)$.

Let χ be sufficiently large and let X be an arbitrary element of $[\mathcal{H}_\chi]^{\aleph_1}$. We show that there is $M \in \mathcal{M}_\chi^\square$ such that $X \subseteq M$ and $P \cap M \leq_\sigma P$.

Fix a well-ordering $<^*$ of \mathcal{H}_χ of order type $|\mathcal{H}_\chi|$. Let $\mathcal{C} = \{C_\alpha : \alpha \in \text{Lim}(\omega_2)\}$ be a \square_{ω_1} -sequence.

Let $(M_\alpha)_{\alpha < \omega_2}$ and $(a_{\alpha, \gamma})_{\alpha < \omega_2, \gamma < \omega_1}$ be sequences defined inductively so that they satisfy the following conditions:

- (0) $(M_\alpha)_{\alpha < \omega_2}$ is a continuously increasing sequence of elementary submodels of $(\mathcal{H}_\chi, \in, <^*)$ of cardinality \aleph_1 .
- (1) $\omega_1, X \subseteq M_0, P, \mathcal{C} \in M_0$.
- (2) For all $\alpha < \omega_2$, $(a_{\alpha, \gamma})_{\gamma < \omega_1}$ is an enumeration of M_α .
- (3) For all $\beta < \omega_2$ we have $(M_\alpha)_{\alpha \leq \beta}, <^* \cap (\bigcup_{\alpha \leq \beta} M_\alpha)^2, (a_{\alpha, \gamma})_{\alpha \leq \beta, \gamma < \omega_1} \in M_{\beta+1}$.
- (4) For all $\alpha < \omega_2$, $P \cap M_{\alpha+1} \leq_\sigma P$.

Let $M = \bigcup_{\alpha < \omega_2} M_\alpha$. Then $P \cap M \leq_{\aleph_2} P$ by the \aleph_2 -IDP of P . From (4) it follows that $P \cap M \leq_\sigma P$. Hence by Lemma 8.15, $\text{SEP}(P \cap M)$. It follows from Lemma

6.3 that there is $\alpha^* \in E_{\omega_1}^{\omega_2}$ such that $P \cap M_{\alpha^*} \leq_\sigma P \cap M \leq_\sigma P$. Since $X \subseteq M_{\alpha^*}$ by (1), the proof is complete if we can show the following:

Claim. $M_{\alpha^*} \in \mathcal{M}_\chi^\square$.

Let $C = C_{\alpha^*}$. C is a cofinal subset of α^* of order type ω_1 . Let $(\xi_\alpha)_{\alpha < \omega_1}$ be strictly increasing enumeration of C . For each limit ordinal $\alpha < \omega_1$ there is $\beta < \alpha^*$ such that $\xi_\alpha \in M_\beta$. Since $C_{\xi_\alpha} = \{\xi_\gamma : \gamma < \alpha\}$ by coherence and since $C_{\xi_\alpha} \in M_\beta$, we have $\{\xi_\gamma : \gamma < \alpha\} \in M_\beta \subseteq M_{\alpha^*}$. Hence

(*) for all $\alpha < \omega_1$, $\{\xi_\gamma : \gamma < \alpha\} \in M_{\alpha^*}$.

Let $\varphi : \omega_1 \rightarrow \omega_1 \times \omega_1$; $\alpha \mapsto (\varphi_0(\alpha), \varphi_1(\alpha))$ be a surjection such that $\varphi \in M_0$.

We now define a continuously increasing sequence $(N_\alpha)_{\alpha < \omega_1}$ of countable elementary submodels of M_{α^*} such that

- (5) $a_{\xi_{\varphi_0(\alpha)}, \varphi_1(\alpha)}, (N_\beta)_{\beta \leq \alpha} \in N_{\alpha+1}$ for all $\alpha < \omega_1$;
- (6) $N_{\alpha+1}$ is the countable elementary submodel of M_{α^*} with $N_{\alpha+1} \in M_{\alpha^*}$ which is minimal with respect to $<^*$ satisfying (5).

That this construction is possible can be seen as follows: By (*) and since the predicate “ $N_\alpha \prec M_{\alpha^*}$ ” can be replaced by “ $N_\alpha \prec M_\eta$ ” for a sufficiently large ordinal $\eta < \alpha^*$, each initial segment of $(N_\alpha)_{\alpha < \omega_1}$ is definable in \mathcal{H}_χ with parameters in M_{α^*} and hence is an element of M_{α^*} .

By (5) $\bigcup_{\alpha < \omega_1} N_\alpha = M_{\alpha^*}$ and $(N_\beta)_{\beta \leq \alpha} \in N_{\alpha+1}$ for all $\alpha < \omega_1$. From Lemma 8.9 it follows that $M_{\alpha^*} \in \mathcal{M}_\chi^\square$. This finishes the proof of the claim and hence of the theorem. \square

Corollary 8.16. *Suppose that \square_{ω_1} holds and $\mathcal{P}(\omega)$ has the \aleph_2 -IDP (in particular this is the case if $2^{\aleph_0} = \aleph_2$). Then $\text{SEP}(\mathcal{P}(\omega))$ implies $\mathfrak{a} = \aleph_1$.*

Proof. Under the assumptions, if $\text{SEP}(\mathcal{P}(\omega))$ then we have $\text{SEP}^+(\mathcal{P}(\omega))$ by Theorem 8.14. Hence $\mathfrak{a} = \aleph_1$ by Theorem 8.12. \square

9. CONCLUSION

As we have mentioned in the introduction, large cardinals are necessary to construct a Boolean algebra A with the IDP but without the WFN. In this sense, IDP and WFN are pretty much the same and it is not surprising that all the interesting set theoretic consequences of $\text{WFN}(\mathcal{P}(\omega))$ that have been discovered so far already follow from $\text{IDP}(\mathcal{P}(\omega))$. Looking at the proofs of the known consequences of $\text{WFN}(\mathcal{P}(\omega))$ or $\text{IDP}(\mathcal{P}(\omega))$, it turns out that most of the time $\text{SEP}(\mathcal{P}(\omega))$ is enough to derive these consequences. An exception could be the equality $\mathfrak{a} = \aleph_1$, which is not known to follow from $\text{SEP}(\mathcal{P}(\omega))$, but which follows from $\text{IDP}(\mathcal{P}(\omega))$. The natural open question is whether $\text{SEP}(\mathcal{P}(\omega)) + \mathfrak{a} > \aleph_1$ is consistent.

One nice feature of $\text{SEP}(\mathcal{P}(\omega))$ is that it holds in Cohen models. This does not have to be true for $\text{WFN}(\mathcal{P}(\omega))$ (assuming large cardinals). We do not know about $\text{IDP}(\mathcal{P}(\omega))$. As it turns out, $\text{SEP}(\mathcal{P}(\omega))$ is relatively robust under slight changes of the definition. It does not matter whether we demand the existence of a single elementary submodel of \mathcal{H}_χ with certain properties, or of stationarily many, or of cofinally many. Therefore it is interesting to know that the strongest variant of SEP

along these lines, IDP, is strictly stronger than SEP. In some sense, we get the best possible result here. There is (in ZFC) a Boolean algebra with SEP but without IDP and it is consistent that $\mathcal{P}(\omega)$ itself is an example.

REFERENCES

- [1] A. Blass, *Combinatorial Cardinal Characteristics of the Continuum*, in Handbook of Set Theory, to appear.
- [2] A. Dow, K.P. Hart, *Applications of another characterization of $\beta\mathbb{N} \setminus \mathbb{N}$* , Topology and its Applications, 122, 1-2, 105–133 (2002)
- [3] M. Foreman, M. Magidor, and S. Shelah, Martin’s Maximum, saturated ideals and non-regular ultrafilters. Part I. AM 127, 1–47 (1988)
- [4] S. Fuchino, S. Geschke, L. Soukup, *The weak Freese-Nation property of $\mathcal{P}(\omega)$* , Archive of Mathematical Logic 40, No.6, 425–435 (2001)
- [5] S. Fuchino, S. Geschke, S. Shelah, L. Soukup, *On the weak Freese-Nation property of complete Boolean algebras*, Annals of Pure and Applied Logic 110, No.1-3, 89–105 (2001)
- [6] S. Fuchino, S. Koppelberg, S. Shelah, *Partial orderings with the weak Freese-Nation property*, Annals of Pure and Applied Logic, 80 (1996)
- [7] S. Fuchino, L. Soukup, *More set theory around the weak Freese-Nation property*, Fundamenta Mathematicae 154, 159–176 (1997)
- [8] I. Juhász, K. Kunen, *The Power Set of ω , Elementary submodels and weakenings of CH*, Fundamenta Mathematicae 170, 257–265 (2001)
- [9] I. Juhász, L. Soukup, Z. Szentmiklossy, *Combinatorial principles from adding Cohen reals*, in Proceedings of Logic Colloquium (1995)
- [10] S. Shelah and J. Steprāns, Non-trivial automorphisms of $\beta\mathbb{N} \setminus \mathbb{N}$ without the continuum hypothesis, Fundamenta Mathematicae 132, 135–141 (1989)

(Fuchino) DEPT. OF NATURAL SCIENCE AND MATHEMATICS, COLLEGE OF ENGINEERING, CHUBU UNIVERSITY, KASUGAI AICHI 487-8501 JAPAN

E-mail address: fuchino@isc.chubu.ac.jp

(Geschke) II. MATHEMATISCHES INSTITUT, FREIE UNIVERSITÄT BERLIN, ARNIMALLEE 3, 14195 BERLIN, GERMANY

E-mail address: geschke@math.fu-berlin.de